BREATHE the lung association

Flu, Pneumonia and COVID-19 at a Glance

What is Influenza (the Flu)?

The flu is a highly contagious illness caused by influenza virus. The influenza virus causes infections of the nose, throat, and sometimes lungs. It spreads easily by coughing, sneezing and direct contact with spit, phlegm or runny nose.

Symptoms

Flu symptoms usually come on quickly. It is important to stay home to avoid making others ill if you experience any of these symptoms.

NOTE

Contact your doctor or other healthcare provider if you believe you have the flu. To limit the spread of disease during the pandemic, it is important to contact your healthcare provider first by phone or online to make an appointment.

What is Pneumonia?

Pneumonia is swelling (inflammation) in one or both lungs that is usually caused by an infection. Many different germs can cause pneumonia, including bacteria, fungi and viruses (including influenza and COVID-19).

Pneumonia can be lifethreatening. It's a leading cause of death and hospitalization among seniors and in people with longterm (chronic) diseases.

Symptoms

The signs and symptoms of pneumonia can range from mild to severe. It depends on your age, if you have other illnesses and what type of pneumonia you have.

What is COVID-19?

COVID-19 is an infectious disease caused by the novel coronavirus. Other types of coronavirus have caused diseases such as MERS and SARS. This respiratory illness is most often spread from person to person via droplets from the nose or mouth. It is also possible to get COVID-19 by coming into contact with surfaces that have the virus present.

Symptoms

Most people who are infected with COVID-19 will have mild symptoms, but some people will have more severe symptoms.

See next page for a list of symptoms.

What are the Symptoms and When to Seek Medical Help


Symptoms	Flu	Pneumonia	COVID-19
Fever	Common	Common	Common
Cough	Common	Common	Common
Shortness of Breath	Possible/severe	Common	Common
Difficulty Breathing	Possible/severe	Unlikely	Common
Chills	Common	Unlikely	Possible
Fatigue/Weakness	Common	Common	Common
Muscle/Body Aches	Common	Unlikely	Possible
Loss of Taste/Smell	Unlikely	Unlikely	Possible
Headache	Unlikely	Unlikely	Possible
Stomach Ache/Diarrhea/Vomiting	Possible/severe	Unlikely	Possible
Sore Throat	Common	Unlikely	Possible
Loss of Appetite	Common	Unlikely	Unlikely
Runny Nose/Sneezing/Watery Eyes	Common	Unlikely	Unlikely
Phlegm/Mucus	Unlikely	Common	Unlikely
Chest Pain	Possible/severe	Common	Possible
Conjunctivitis	Unlikely	Unlikely	Possible/rare
Rash on Skin	Unlikely	Unlikely	Possible/rare
Discolouration of Fingers/Toes	Unlikely	Unlikely	Possible/rare

There are also some severe flu symptoms in children and adults.

If you experience any of these, seek immediate medical attention.

Symptoms	Child	Adult
Fast breathing, difficulty breathing, shortness of breath		
Blue or grey skin colour		
Pain or pressure in chest or stomach		
Sudden dizziness		
Blue or grey lips		
Confusion		
Dehydration (not peeing)		
Unable to stop vomiting (throwing up)		
Not waking up, not paying attention to anything		
Crankiness		
Seizures		
Symptoms improve but then come back; cough is worse, fever		

Symptoms at a Glance for Adults



How Can You Prevent Getting Sick?

Getting the annual flu shot, proper hand-washing and staying home when sick are all key in making sure you are as healthy as possible. The flu shot will not protect against COVID-19, but it will help reduce your risk of getting the flu and COVID-19 at the same time.



Take These Preventative Measures	Flu	Pneumonia	COVID-19
Get vaccinated.			
Wash your hands frequently and thoroughly for at least 20 seconds with warm water and soap.	•	•	•
Avoid touching your eyes, mouth and nose.			
Stay away from others when you or when they are sick.			
Stay home from work, school or public places when sick.			
Cover your mouth and nose if you cough or sneeze with your elbow or a tissue.	•		•
Throw away used tissues and wash your hands immediately.			
Avoid smoking. People who smoke are at a higher risk of pneumonia, as are children of parents who smoke.			
Smoking any kind of tobacco reduces lung capacity and increases the risk of many respiratory infections and can increase the severity of respiratory diseases.		•	•
Do not share cups/eating utensils with people who are sick.			
Maintain distance of at least 6 feet (2 meters) from others.			
Wear an appropriately fitted face mask.			



Fact Sheet | Flu, Pneumonia and COVID-19 at a Glance

What are the Treatments?

Most people with mild cases of either of the diseases can manage their conditions at home.

Treatments	Flu	Pneumonia	COVID-19
Stay at home			
Drink lots of fluids			
Rest			
Antibiotics (* for bacterial pneumonia only)		•*	
Over the-counter medication to help with headache, muscle aches and fever such as acetaminophen (for example, Tylenol [®]) or ibuprofen (for example, Advil [®])	•		•
Eat a nutritious diet			

The Flu

Children and teenagers younger than 18 years of age who have chicken pox, cold or flu symptoms should not be given ASA (acetylsalicylic acid) products (for example, Aspirin[®]) before a doctor or pharmacist is consulted. Reye's Syndrome, a rare and potentially fatal illness is reported to be associated with ASA when used to treat chicken pox, cold or flu symptoms in children.¹ If you're not sure what to take, consult a doctor or pharmacist.

Take antiviral medication, if prescribed. Not everyone will be given antivirals because it is only recommended for people with severe illness, people who require hospitalization and people who are at high risk of complications.

Note: If you have asthma, follow the advice in your asthma action plan. If you have COPD, follow the advice in your COPD action plan. You may have to take extra medicine to control your symptoms. If it's hard to breathe or your rescue medicine isn't working, call 911 or your local emergency number.

Pneumonia

Some people with severe pneumonia will need to go to the hospital to have additional support of oxygen and medication intravenously. If you are prescribed antibiotics it is important to take the medicine as your doctor prescribes, even if you start to feel better. Do not take cough or cold medicine unless your doctor says it is okay.

If you have a milder pneumonia, you will usually begin to feel better within a few days or a week. If you have severe pneumonia, where you spend time in the hospital you may take a number of weeks to feel better.

People who have been admitted to the hospital with other medical conditions and develop pneumonia may become very ill and could require stronger antibiotics intravenously (IV).

Note: Antibiotics do not treat viral pneumonia. Some of the treatment of viral pneumonia may include drinking lots of fluids, corticosteroid medicine (to reduce inflammation), oxygen therapy and rest.

COVID-19

Currently, there are only treatments for COVID-19 available for those with severe disease who are in hospital. Clinical trials are underway to find both preventative and treatment medication for COVID-19. If you are concerned about your symptoms, contact your doctor.²

¹ https://www.healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2013/36303a-eng.php

² https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/symptoms.html